

1 Suspect Screening and Non-Targeted Analysis of Drinking Water Using
2 Point-Of-Use Filters

3 Supporting Information

4 Text and Figures

5 Seth R. Newton¹, Rebecca L. McMahan², Jon R. Sobus¹, Kamel Mansouri^{2,3}, Antony J. Williams³,
6 Andrew D. McEachran^{2,3}, and Mark J. Strynar¹

7 1. United States Environmental Protection Agency, National Exposure Research
8 Laboratory, Research Triangle Park, NC 27709, United States.

9 2. Oak Ridge Institute for Science and Education Research Participant, 109 T.W. Alexander
10 Drive, Research Triangle Park, NC 27709, United States.

11 3. United States Environmental Protection Agency, National Center for Computational
12 Toxicology, Research Triangle Park, NC 27709, United States.

13

14

15

16 **Materials**

17 Dichloromethane (DCM) and methanol (B&J Brand High Purity Solvent) were purchased from
18 Honeywell Burdick & Jackson (Muskegon, MI, USA). Ammonium acetate and ammonium
19 formate were purchased from Sigma Aldrich (St. Louis, MO, USA). Ultrapure deionized (DI)
20 water was generated in house from a Barnsted Easypure UV/UF (Dubuque, IA, USA) coupled
21 with activated charcoal and ion exchange resin canisters.

22

Figure S1 - Bubble plot of all features found in the Brita[®] extract colored by positive and negative ionization mode, and whether they were matched or unmatched to a formula in the database. The size of the bubble is proportional to the size of its corresponding chromatographic peak.




Figure S2 – PCA using the parsed formulas from the water filters and house dust samples from Rager *et al.* C = carbon, Cl = chlorine, F = fluorine, H = hydrogen, N = nitrogen, O = oxygen, P = phosphorous, RT = retention time, S = sulfur. Bromine was omitted due to its lack of occurrence (one house dust formula and no Brita® formulas contained bromine).




Figure S3 – Loadings plot from the PCA of product-use categories.




Figure S4 – A) Chromatograms of decarboxylated PFOA, B) Chromatogram of PFOA, and C) Spectrum of the two compounds showing the mass difference between the two (43.9901) corresponding to CO₂




Figure S5 – Chromatograms of A) m/z 518.8796 unknown peak in Pittsboro Tap sample, B) m/z 564.8848 unknown peak in Pittsboro Tap sample, C) m/z 564.8848 unknown peak in a method blank, and D) blank subtracted spectrum at m/z 564.8848 in Pittsboro Tap sample with theoretical spectrum of the formula $C_{13}H_{22}Cl_7O_7P$ ($C_{12}H_{20}Cl_7O_5P$ + formate adduct) in red boxes.

